

CAHOW RECOVERY PROGRAM 2010 – 2011 Breeding Season Report



**Compiled by: Jeremy Madeiros, Conservation Officer
Terrestrial Conservation Division
Department of Conservation Services
Ministry of Public Works**

BERMUDA GOVERNMENT



Fig. 1: Two-week old Cahow chick removed from burrow on nesting island

DEPARTMENT OF CONSERVATION SERVICES

**Applied Ecology Section
Terrestrial Conservation Division**

BERMUDA

RECOVERY PROGRAM FOR THE CAHOW

(BERMUDA PETREL)

Pterodroma cahow

Breeding Season Report

For the Nesting Season October 2010 to June 2011

CONTENTS:

Page No.

(1) Executive Summary:	5
(2) Objectives of Cahow Recovery Program:	7
(3) Introduction:	8
(4) Cahow Recovery Program – Review of 2010/2011 Nesting Season:	11
(5) Summary of 2010 / 2011 Nesting Season:	14
(6) Breakdown of Breeding Season Results by Nesting Island:	15
(7) Progress Report on New nesting Colony on Nonsuch Island:	16
(8) Update on Cahow Banding Program 2011:	20
(9) Update on Oceanic Range Study Using Archival Geolocational Loggers:	22
(10) Planned Future Management Actions and Research:	23
(11) Acknowledgements:	24

List of Figures

Cover photo: Adult Cahow flying over Nonsuch Island at night (Chris Burville)

Fig. 1: Two-week old cahow chick removed from nest burrow	(2)
Fig. 2: Half-fledged downy Cahow chick on Nonsuch Island	(6)
Fig. 3: Location of original nesting islands and new Nonsuch colony	(9)
Fig. 4: Chart showing no. of breeding pairs over 50-year period.....	(12)
Fig. 5: Cahow nest burrows on Nonsuch Island 2011	(16)
Fig. 6: Cahow chick on Nonsuch Island out to exercise at night	(18)
Fig. 7: J. Madeiros and D. Wingate with Cahow chicks, May 2011	(19)
Fig. 8: Identification band fitted on leg of adult Cahow	(20)
Fig. 9: Oceanic range of Cahow with all data points from loggers	(22)
Fig. 10: Travels of 1 adult Cahow over 12 months as recorded by logger..	(23)
Fig. 11: Adult Cahow in Nest with Large Egg (Jan 2011)	(26)

- Fig. 4: Leila Madeiros; Figs. 5, 9, 10: Mandy Shailer;
- Fig.7: Kathryn Deane. All other photos: Jeremy Madeiros

(1): EXECUTIVE SUMMARY:

Key Words: Cahow, Banding, Breeding Success, Nonsuch Island, Social Attraction, Translocation.

The Cahow Recovery Program is a long-term management, research and recovery program for Bermuda's National Bird, the critically endangered Cahow, or Bermuda petrel. This program is focused on enabling the increase of the breeding population through the control or elimination of threats to the species, provision of additional artificial nesting burrows, and the establishment of entirely new nesting colonies. The last is underway using translocation of near-fledged chicks from their original nesting islets to artificial burrows at a new colony site on Nonsuch Island.

The Cahow, which nests only on the Islands of Bermuda and was thought to have originally numbered more than half a million birds, was catastrophically affected by the arrival of humans on the island in the early 1600s. This was due both to direct hunting by the settlers and by invasive predators introduced by man, such as Rats, Cats, Dogs and Pigs. After less than 20 years of settlement, the Cahow had declined to the point where it was thought to be extinct, a belief that persisted for over 350 years until the rediscovery in 1951 of a tiny remnant population of breeding pairs on four tiny offshore islets.

Since 1960, a conservation and recovery program has been in place that has addressed and controlled most threats to the species. This program was administered by Dr. David Wingate until his retirement in 2000, since which it has been administered by the author of this report. This program has enabled the breeding population to begin a slow, but accelerating increase from only 18 pairs producing a combined 8 chicks annually in the 1960s to a new record number of 98 breeding pairs in 2011, producing a new record total of 56 fledged chicks. Increased knowledge and public interest in the Cahow has been brought about from several films and documentaries that have been completed highlighting the conservation work being carried out on the species.

The main threats to the Cahow include the erosion and flooding of the present nesting islets by storm activity and continuing sea-level rise, the threat posed by Rats and other invasive species swimming to these islets, a lack of sufficient numbers of potential nest burrows or rock crevices, and nest-site competition with the Longtail or White-tailed Tropicbird *Phaethon lepturus catsbyii*.

Following are some of the highlights for the 2011 Cahow nesting season:

- The new nesting colony of Cahows established on Nonsuch Island by the translocation of chicks between 2004 and 2008 continues to grow, with 7 pairs established in nest burrows and laying eggs. From these, four chicks hatched and successfully fledged out to sea. New pairs and prospecting activity was noted in 4 additional nests, and a total of 21 of the translocated birds have so far returned to Nonsuch as adults, in addition to 2 non-translocated Cahows attracted to the new colony by the returned translocated birds.

- The total breeding population of the Cahow is approaching 100 pairs, producing nearly 60 fledged chicks, numbers which almost certainly have not been seen since the early 1600s.
- Despite the impact of large, powerful hurricane ‘Igor’ in September, 2010, all damage and erosion to nesting burrows was repaired in time for the new nesting season, and there appears to have been little or no effect on breeding success in the 2010-2011 breeding season.
- The second year of deployment of archival geolocational data loggers was completed with at least four additional tags being recovered, at least two of which were deployed on individual Cahows for two full seasons. The data from these tags will, when fully analyzed, be used to establish the at-sea range of the Cahow, including favored foraging areas and migration routes.

The Recovery Program continues to achieve its primary objective of increasing the Cahow breeding population and the number of successfully fledged chicks. Increasing public and scientific interest in the Cahow and the Recovery Program has been helped by the production of a 30-minute documentary, titled “Higher Ground” by LookTV Bermuda and a segment of the “Learn-alots” TV show produced by Panatel Ltd. focused on the Cahow Recovery Project. In addition, two scientific papers on the Cahow produced jointly by the Government Conservation Officer and scientists from the New South Wales Department of the Environment and Climate Change have now been accepted for publication by Conservation International journal. These papers deal with Cahow breeding biology and with the Cahow translocation project.

Full details on the 2010 – 2011 breeding season are given in the following report, in addition to research and management proposals for the next several years.



Fig. 2: Half-fledged downy Cahow chick on Nonsuch Island

(2): Objectives of Cahow Recovery Program:

The Cahow Recovery Program was set up to co-ordinate management and research efforts for the Bermuda petrel or Cahow *Pterodroma cahow*, which is recognized as one of the rarest seabirds on earth. The Cahow is endemic to the islands of Bermuda, nesting no-where else of earth and numbering less than a hundred pairs.

This program has a number of short and long-term objectives, which are covered in full in the Cahow Recovery Plan (Madeiros, 2005); briefly put, these objectives are as follows:

- (1) To prevent nest-site competition with the White-tailed Tropicbird (*Phaethon lepturus catsbyii*) through the use of wood “baffle plates” at the entrances of all Cahow nest burrows to prevent Tropicbirds from entering.
- (2) To regularly monitor all nesting and nearby islands for the presence of rats (*Rattus rattus* and *R. norvegicus*) and, when their presence is detected or suspected, to eradicate them by use of anticoagulant rodenticides, using bait and boxes provided by or purchased from the Health Department.
- (3) To continue the program of construction of additional nest burrows at all appropriate nesting islands, and at locations where new nesting colonies are being established, to support a continued increase in the breeding population.
- (4) To establish new nesting colonies of Cahows on larger, more elevated islands free of mammal predators, and which are safer from hurricane erosion and have the potential of supporting larger populations of the birds; this is already being achieved on Nonsuch Island and is planned for the future on Southampton Island.
- (5) To learn more about the biology of the species through an ongoing banding program initiated in 2002; also through developmental studies of Cahow chicks and morphometric measurements of adult Cahows.
- (6) To carry out studies of the oceanic range of Cahows using archival geolocational data loggers attached to individual birds, which record daily position fixes for periods of up to 2 years. These have already been successful in recording foraging areas and migration routes used by Cahows, both when they forage for food for their chicks during the breeding season, and where they travel during the summer, non-breeding season.

(3): Introduction:

The 2010-2011 Nesting Season of Bermuda's national bird, the endemic and critically endangered Bermuda Petrel or Cahow (*Pterodroma cahow*) began in late October 2010 and ended in mid-June 2011 with the departure of the last fledglings out to the open ocean from their nesting burrows. This nesting season has been marked by the continued increase in the numbers of breeding Cahows, with numbers of breeding pairs getting close to the 100 pair mark. A record number of 98 pairs nested on the original nesting islets and at the new colony site on Nonsuch Island, with a total of 56 chicks successfully fledging out to sea, also a new record high number. These are up from a total of just 18 nesting pairs and 8 fledged chicks when the recovery program began in 1960.

The Cahow is subject to pressure from a number of threats and limiting factors, including nest site competition from the native White-tailed Tropicbird, the threat of introduced mammal predators (especially rats) swimming out to the nesting islets, lack of suitable nesting sites at the present islets, and the annual threat of massive erosion and damage to the islets from hurricane waves and storm surge. Much of the present management carried out on the Cahow through the Recovery Program has been focused on addressing and overcoming these various threats to the species, with a high degree of success.

The single most important threat to the Cahow is now considered to be ongoing damage and major erosion to the nesting islets from severe hurricanes and storms, coupled with accelerating sea-level rise. After a thirty-five year period (early 1950s to late 1980s) with few to no major impacts from hurricane waves and storm surge, there have no fewer than nine major hurricane impacts to the nesting islets from hurricanes between 1989 and 2011 (hurricanes "Dean" in 1989; "Lily" in 1991; "Felix" in 1995; "Gert" in 1999; "Fabian" in 2003, "Florence" in 2006; "Bill" in 2009; "Igor" in 2010; and "Katia" in 2011).

The impacts from these hurricanes have included the undermining and collapse of large sections of the islands, and breaking away of large chunks of limestone and cliff face, coupled with overwashing of the smaller islets, when large waves break completely over the island. This has resulted in both complete destruction of many of the nesting burrows used by Cahows, and in damage to and filling in by rocks and debris of much of the remainder. This has resulted in much dangerous and labor-intensive work needed to repair nests at the end of hurricane season in many years before the Cahows return for their nesting season. It is important to repair as many of the original nests as possible, as their loss will result in long-established pairs breaking up and needing several years to re-establish new nests with new mates.

To address this threat, a major component of the recovery program is to establish new nesting colonies on islands that are both larger and more elevated than the original tiny nesting islets, which are generally only half an acre (0.2 Ha) in area each (See Fig. 3). This has already been accomplished on Nonsuch Island, which at 16.5 acres is the largest isolated island in the Castle Harbour area, and is maintained rat-free as part of the Living Museum ecological restoration project. A new colony has been established with 12 active

nest burrows occupied by returned Cahows which had originally been translocated as chicks from the original nesting islands to a new complex of artificial burrows on Nonsuch Island. 105 chicks were translocated between 2004 and 2008, and were then hand-fed and measured until they fledged out to sea, imprinting on Nonsuch Island in the process. By 2011, a total of 28 of the translocated Cahows had returned as adults, of which 21 so far have been recaptured in nests back on Nonsuch Island.

As this first effort to establish a new Cahow nesting colony is proving to be successful, plans are now underway to establish a second colony site elsewhere on Nonsuch Island, followed by an attempt to establish a new colony on Southampton Island. Although smaller than Nonsuch at 2.36 acres, Southampton Island is still larger than the original nesting islets and is located only 100 meters from a current Cahow nesting colony. As at the first Nonsuch colony site, these additional colony establishment efforts will be undertaken using a combination of translocation of chicks from the original islets, and sound attraction, using pre-recorded courtship calls played back at the new colony sites to attract returning Cahows.

In addition to the encouraging return of translocated Cahows and the continued growth of the new nesting colony on Nonsuch Island, new pairs of nesting Cahows established in nest burrows on three of the original nesting islets (Inner Pear Rock, Horn Rock and Green Island). As a counterpoint, two long-established pairs of Cahows appeared to break up with one bird from each pair not returning for the nesting season, although this is probably due to normal natural mortality and well within natural mortality rates in the growing Cahow nesting population.

The final months of filming for a new 30-minute documentary on the Cahow Recovery Project by local television channel LookTV (channel 1) were shot during this nesting season and the finished show first aired on local cable television in June, 2011. This documentary, called "Higher Ground", was filmed over five years and focuses on both the translocation project to establish a new nesting colony on Nonsuch Island and the hurricane erosion and sea-level rise that makes this project imperative for the continued recovery of the species. It also records the return of translocated birds to the Nonsuch site as adults, the formation of nesting pairs in artificial nest burrows on the island, and the hatching of "Somers", the first Cahow chick to naturally hatch on Nonsuch since the 1620s (See 2009 Cahow Nesting Season Report for full details).

Following are highlights from the 2010 – 2011 Cahow nesting season:

- (1) The total number of active nesting pairs of Cahows increased to a record high of 98, compared to 92 nesting pairs in the 2009 – 2010 nesting season and 55 pairs in the 2000 – 2001 seasons.
- (2) Coupled with the record number of nesting pairs of Cahow, a new record number of 56 chicks successfully fledged from the nesting islands, compared to 52 chicks that were fledged in the 2009 – 2010 nesting season.
- (3) The new nesting colony on Nonsuch Island, which was established by the translocation of near-fledged chicks and sound attraction techniques between 2004 and 2009, has continued to grow. As of June 2011, a total of 21 of the translocated birds have returned to Nonsuch Island as adults, with 7 nesting pairs laying eggs in 2011. Of these, four chicks hatched and fledged successfully out to sea, while new pairs or prospecting adult Cahows have established themselves in five additional nests. This means that a total of 12 nest burrows on Nonsuch Island now have nesting or pre-nesting activity.
- (4) The second year of deployment of geolocational archival data loggers was completed with at least 4 additional loggers being retrieved from adult Cahows that they had been attached to. At least two of these loggers had been deployed on Cahows for 2 full years. Data from these tags will be analyzed and used to help establish the oceanic range of the Cahow, including favored foraging areas and migratory routes. The data from the first year of deployment of these tags has revolutionized understanding of the distances that Cahows fly when collecting food for their chicks, where they go over the summer, non-nesting season, and the total distances travelled by Cahows over periods spanning up to a year or more.

(4): Cahow Recovery Program – Review of 2010/2011 Nesting Season and Management Actions:

The first management actions of the 2010 – 2011 Cahow nesting season were necessary repairs to a number of nesting burrows that had been damaged by high seas during hurricane “Igor”, which passed over the Bermuda area during the 19th and 20th September 2010. This hurricane was unusually large and hit Bermuda with winds gusting from 85 to 110 mph, with waves up to 25’ partly submerging three of the nesting Islands, and causing moderate to severe erosion to all four islands. 11 of the heavy concrete nest lids that cover the Cahow burrows were swept overboard by waves breaking over Long Rock, and 8 nest lids from Green Island, with many of the nests also being filled by rocks and debris. On the 24th September, Mr. Peter Drew was able to dive up most of these nest lids from where they had been swept off the islands. Further work in clearing rocks, vegetation and debris from the nests and their entrances were completed in time for the arrival of the first Cahows back to the nesting grounds. Rat poison was also set out on the nesting islands to ensure that they were rat-free for the coming season.

During the first week of October, all of the Cahow nesting burrows were unblocked and the entrance Baffles which prevent nest invasions by White-tailed Tropicbirds during the spring period were removed and taken to Nonsuch Island for storage. The first Cahows were confirmed as returning to their nest burrows by the 18th October, 2010, officially starting the nesting season. Most of the active nesting pairs had returned to their nest burrows by the 31st October.

Grass was collected in trash bags from Nonsuch Island and transported out to the smaller nesting islets by early November to provide sufficient material for the Cahows to be able to build their nests. This is necessary as these islets are so rocky that they have insufficient grass and vegetation for Cahows to build good nests to cushion the eggs and reduce accidental breakage. From the 20th October until the beginning of December, visits to the nesting islands to check all Cahow burrows were carried out on average 4 to 5 days a week, as weather conditions allowed. During these checks, any Cahows found were checked for band numbers, weight and body condition. This period is when breeding pairs carry out courtship activity and build a nest in their burrows from vegetation gathered outside the burrow entrance.

During most of the month of December, 2010, the Cahow breeding population returned out to sea, giving a break during which some modifications and repairs were carried out to a number of the nest burrows damaged during hurricane “Igor”. This period, which is termed the pre-egg laying exodus, occurs because both partners of a nesting pair need to carry out a 5-week period of intensive feeding, the female to develop her single large egg and the male to accumulate ample fat reserves for the first long shift of egg incubation, lasting up to 2 weeks with no food, while the female returns to sea to recover after laying an egg which can mass between one-quarter and one-fifth of her total body weight.

The first Cahows returned to the nesting islands from their pre-egg laying exodus at the beginning of January, 2011, with the first eggs being confirmed on the 5th January (See Fig. 5). The egg incubation period, which lasts about 53 days, is the main period in which incubating adults are checked to determine sex and band numbers to determine which birds are returning to which nests. Checks to the nesting islands were made impossible at times by frequent gale-force winds as winter storm systems passed over the Bermuda area; one low pressure system on the 23rd January produced winds gusting to 69 mph.




Fig. 4: Chart showing number of breeding pairs and fledged young Cahow over 50-year period

The first Cahow chick was confirmed hatching on the 23rd February, with more by the 28th February. By the 27th March, a total of 59 chicks had hatched, some of which had to be confirmed from the deeper natural nests by the use of infra-red “burrow-scope” equipment purchased during the previous year for just that purpose. Once the chicks had hatched, a sub-sample of about 30 chicks was chosen which were then checked at least twice weekly, weather conditions permitting, for weight, wing chord length, and plumage development. This information is being used for a study of chick growth rates, but also has practical application in identifying chicks that are neglected or abandoned prematurely by the adults and need to be taken into care. Most importantly, it is also used for identifying when chicks that are being translocated are at the optimal age to be moved to their new nest sites. These

checks of Cahow chick growth rates continue until the chicks depart to sea between mid-May and early June.

The chicks are fitted with identification bands on their left legs once their adult plumage covers more than half their body, usually at 70 days of age or older (adult birds whose ages are not known have their bands fitted to their right legs). During the 2011 Cahow nesting season, a total of 32 chicks were fitted with identification bands, out of a total of 56 chicks which successfully fledged (57% of all chicks). As of June, 2011, a total of 408 Cahows have now been fitted with identification bands, the majority of which have been fledgling birds. The return and eventual nesting of these known-age birds has already given a wealth of previously unknown information on the age of first return of young adult Cahows, time of new pair establishment and first successful production of chicks, and nest site and pair fidelity (faithfulness).

Several Cahow chicks died at various points in their development, some from parental abandonment and others from unknown causes. This is a normal occurrence during a breeding season, although all efforts are made to save as many abandoned chicks as possible (See following paragraph). Despite this, a record high number of chicks fledged successfully out to sea. The first Cahow chicks fledged out to sea on May 21st, from Inner Pear Rock, while the last chick did not depart until 28th June, 2011.

Three Cahow chicks were abandoned prematurely by their parents, and had to be taken into care to feed them and enable them to accumulate enough fat reserves to fledge successfully out to sea. These chicks were moved from the C4 and C27 nest burrows on Horn Rock, and the D4 burrow on Inner Pear Rock, to empty artificial nest burrows at the translocation site on Nonsuch Island. Once under care, they were fed daily on fresh Anchovies and fresh unfrozen squid, using the same techniques developed for the translocation project to establish the new nesting colony on Nonsuch. Two of the chicks from the Horn Rock C4 nest and the IPO D4 nest recovered, gained weight and eventually matured and fledged successfully out to sea from Nonsuch Island. The third chick, from the Horn Rock C27 nest began to regurgitate all food that was fed to it, sickened and eventually died, and was determined to have been too emaciated to have been saved by the time it was brought in.

The newly established Cahow nesting colony on Nonsuch Island had a very successful year and continues to grow in both the number of nesting pairs and the number of successfully fledged chicks. A total of 7 pairs laid eggs in nest burrows, from which 4 chicks hatched and eventually fledged out to sea, while the number of adult Cahows confirmed as returning to Nonsuch Island has exceeded 20 (See section 7 for full details).

(5): Summary of 2011 Cahow Nesting Season:

The 2011 Cahow nesting season has been highlighted by the highest number of successfully fledged chicks to be recorded since the rediscovery of the species, and by the continued growth of the new nesting colony that has been established on the Nonsuch Island Nature Reserve. This new colony also produced 4 successfully fledging chicks.

The numbers of nesting pairs of Cahow have increased to a record high number of 98, of which 56 produced successfully fledging chicks. This represents a breeding success rate of 57.14%. In addition, new prospecting or pre-breeding activity was recorded at 6 additional nest sites, including 3 new nest sites on Nonsuch Island (See Figs. 6 – 10 for maps of nesting islands and status of nest sites).

Following is a summary of the 2011 nesting season results:

Total number of nest burrows with confirmed nesting activity:	98
Number of new nest sites with prospecting activity:	6
Total number of confirmed successfully fledged chicks:	56
Total number of active nest sites with unsuccessful nesting:	42
Number of failures from nest sites with observable nest chambers:	39
Number of failures from nest sites with non-observable nest chambers:	3

Breakdown for causes of breeding failure from nests where observation of nest chambers was possible:

Chick died hatching:	2
Chick died later in development:	4
Eggs broken or pipped:	15
Egg broken during nest check:	1
Non-hatching / infertile eggs:	12
Egg buried or knocked off nest:	2
Egg washed off nest by storm waves:	1
Chick disappeared shortly after hatching – Land Hermit Crab predation?:	2

(6): Breakdown of Breeding Season Results by Nesting Island:

LONG ROCK:

Active nest burrows with nesting confirmed (eggs laid and/or chick hatched):	13
New nest burrow prospected by confirmed pair:	0
Nest burrows with successfully fledged chicks:	9
Nest burrows with confirmed failed nesting:	4

INNER PEAR ROCK:

Active nest burrows with nesting confirmed:	19
New nest burrows prospected by confirmed pairs:	0
Nest burrows with successfully fledged chicks:	12
Nest burrows with confirmed failed nesting:	7

GREEN ISLAND:

Active nest burrows with nesting confirmed:	21
New nest burrows prospected by confirmed pairs:	2
Nest burrows with successfully fledged chicks:	10
Nest burrows with confirmed failed nesting:	11

HORN ROCK:

Active nest burrows with nesting confirmed:	38
New nest burrows prospected by confirmed pairs:	1
Nest burrows with confirmed successfully fledged chicks:	21
Nest burrows with confirmed failed nesting:	17

NONSUCH ISLAND: (See Fig. 5)

Active nest burrows with nesting confirmed:	7
New nest burrows prospected by confirmed pairs:	3
Nest burrows with confirmed successfully fledged chicks:	4
Nest burrows with confirmed failed nesting:	3



Fig. 5: Cahow nest burrows on Nonsuch Island in 2011

(7): Progress Report on New Cahow Nesting Colony on Nonsuch Island:

Between 2004 and 2008 a project was carried out to attempt to establish a new nesting colony of Cahows on a larger, more elevated nesting island than the original nesting locations, which have become increasingly threatened by massive erosion and flooding caused by hurricanes. The Cahow Translocation Project involved the translocation of near-fledged Cahow chicks at approximately 18 days before fledging from their nests on the four original nesting islets to a new complex of artificial burrows constructed on the Nonsuch Island Nature Reserve. Since the original nesting islands range only between 0.5 to 1.0 acres in size, with maximum elevations ranging from 15' (5m) to 32' (10m), Nonsuch Island, which is both much larger at 15.5 acres (6.5 Ha) and of higher elevation at up to 60' (19m) offers much greater protection both from flooding and erosion. It also has a much larger area to allow for growth of the Cahow breeding population to self-sustaining levels, and contains adequate soil and forest cover to eventually allow the Cahows to dig out their own nest burrows.

Gadfly petrels such as the Cahow generally return when mature to the same area that they originally departed from as fledglings, a trait known as *site faithfulness*. To this end, a total of 105 Cahow chicks selected from all four of the original nesting islets were moved to the new nest burrow complex on Nonsuch Island over a five-year period between 2004 and 2008. Once in their new nests on Nonsuch, the translocated chicks were banded and fed daily on imported squid and locally sourced fresh Anchovies and Pilchards, and their weight, wing growth and plumage development recorded daily until they were fully developed. At this point feeding was discontinued, emulating normal parental behavior, and the chicks were then monitored through their exercise period when they emerge nightly to exercise flight muscles and imprint on their surroundings (See Fig. 6), until they finally fledge to the open ocean on their own. A total of 102 translocated Cahow chicks eventually fledged successfully from Nonsuch by 2008.

In addition, a solar-powered sound attraction system was set up at the translocation site to help attract returning birds, encouraging them to land and prospect for new, empty nest burrows. This was undertaken because while it is known that Gadfly petrels tend to return to the original site they fledged from, they also strongly prefer to nest in close proximity to already active nest sites, a trait known as *social attraction*. Until a nucleus of nesting pairs is established on Nonsuch Island, the sound system will broadcast a recording of Cahow courtship calls automatically at night to encourage returning, newly matured translocated birds that an established nesting population is in residence on Nonsuch.

By 2008, the first four returning Cahows translocated three years earlier to Nonsuch were recaptured back at the translocation site on Nonsuch, and their identities confirmed from their band numbers. By the end of that year's nesting season, some of these birds had already been seen prospecting nest burrows at the translocation site.

The 2009 nesting season experienced a great success with the first pairs of Cahows establishing in nest burrows on Nonsuch, culminating in the first egg laid and chick hatched on this Island since at least the 1620s. This chick, known as "Somers" in honor of Admiral Sir George Somers, commander of the "Sea Venture" which was wrecked on reefs off Bermuda in 1609 (a event which led to the permanent settlement of Bermuda three years later), fledged successfully to sea on the 17th June of this year. In addition, a total of 15 returned translocated Cahows were recorded back at the translocation site by the end of this season. Both the Government Minister of the Environment and Sports, Mr. Glenn Blakeley and Acting Minister Mr. Nelson Bascombe travelled out to Nonsuch Island to see the chick, which was the subject of an official Government news release that drew interest from Reuters news and environmental publications around the world.

In the 2009 / 2010 season, the number of established breeding pairs carrying out nesting activity and laying eggs rose to 4, although only one of these produced a successfully fledging chick. This chick, known as "Bermudiana", was produced by the same pair from the R818 nest that had produced the "Somers" chick the previous year. The other three nests that produced unsuccessful eggs were the R831, R832 and R834 burrows. Newly established pairs of Cahows were also confirmed in three additional nests (R816, R817 and R837). The total number of returning birds recaptured on Nonsuch Island that had been

translocated as chicks rose to 17, one from the 2004 translocation cohort, nine from the 2005 cohort, six from the 2006 cohort and one from the 2007 cohort.

For the most recent, 2010 – 2011 nesting season, the new Cahow nesting colony on Nonsuch Island has continued to grow both in numbers of nesting pairs, and in that of successfully fledged chicks. Nesting activity with eggs laid was confirmed in a total of 7 burrows (R816, R817, R818, R831, R832, R834, R837), with chicks hatching in 4 of them (R817, R818, R832, R837). All of these chicks, which were identified by the names “Carter”, “Casper”, “Whitney” and “Lefroy” respectively, fledged successfully out to sea by the 17th June 2011 (See Figs.10 and 11).



Fig. 6: Cahow chick from # 832 nest (Whitney) on Nonsuch Island out to exercise at night, June, 2011

In addition, the number of Cahows translocated to Nonsuch Island as chicks that have returned as adults has risen to 28, representing birds from all five years that chicks were translocated to Nonsuch. Out of these, 21 have been recaptured back on Nonsuch Island itself, while the other seven returned birds have been recaptured back on the original nesting islets. This represents a rate of 75% of the returned, translocated birds that have been recaptured back at their original point of departure (the Nonsuch Island translocation site). It is also very noteworthy that at least two non-translocated Cahows have also been recaptured on Nonsuch Island, one of which has paired up with a translocated bird.

Table 1 shows the breakdown of all translocated Cahows that have been confirmed as returning to the nesting islets as of June, 2011:

TABLE 1: Returns of Cahow Chicks Translocated to Nonsuch Island by June, 2011

YEAR Of chick cohort	Total Returns by 2011	Returns to Nonsuch	Returns to other islets
2004 (14 chicks fledged)	5 (28.5%)	1 (7.1%)	4 (21.4%)
2005 (21 chicks fledged)	12 (57.1%)	9 (43%)	3 (14.1%)
2006 (21 chicks fledged)	7 (28.6%)	7 (28.6%)	
2007 (24 chicks fledged)	3 (12.5%)	3 (12.5%)	
2008 (22 chicks fledged)	1 (4.5%)	1 (4.5%)	
Totals: 102 chicks fledged	Total = 28	Total = 21	Total = 7



Fig. 7: J. Madeiros and D. Wingate on Nonsuch Island with two of the Cahow chicks hatched on Nonsuch during the 2011 nesting season

(8): Update on Cahow Banding Program for 2011:



Fig. 8: Identification band fitted on leg of adult Cahow

An important component of the Cahow Recovery Program has been a program to fit identification bands or rings to the legs of as many Cahows as possible. Banding of Cahows has been carried out since 2002, with 5.5 mm metal bands made of a strong, corrosion-resistant alloy called incoloy being fitted to the right legs of adult Cahows and the left legs of fledgling birds. These bands can last for the life of the birds and are imprinted with a unique code and a return address. The bands used for the Cahows are

made especially for Bermuda by Porzana Ltd. (U.K. Wetlands Trust affiliation) in the United Kingdom.

These bands are vital for allowing easy positive identification of individual Cahows in the field, and enable researchers to follow these birds for essentially their entire breeding lifespan through recaptures over successive years.

The Cahow Banding Program has now been underway for ten years, during which a total of 411 Cahows have been fitted with identification bands as of June, 2011. This includes 115 birds banded as breeding adults, and 296 birds banded as chicks before fledging to sea. In other terms, these numbers represent almost 60% of the current adult breeding population, and about 85% of all chicks to have fledged over the last ten years. It should be noted that this does not represent the total number of living Cahows, as less than half of all fledged chicks survive their first years at sea to return to the breeding islands.

Banding of the adult Cahows gives valuable and previously unknown data on pair faithfulness (the tendency for a mated pair to stay together for many years) and site faithfulness (the tendency for a mated pair to return to the same nest site over an extended period of years).

The banding of Cahow chicks has already provided information on many aspects of behavior, including the following:

- 1) The survival rates of chicks during the period between fledging to sea and their first return to the nesting grounds as sexually mature young adults;
- 2) The age of chicks upon their first return to the nesting grounds, and whether this differs between male and female birds;
- 3) The period of time between the first return of the young birds, to the choosing of nest sites and mates, and the first nesting attempts;
- 4) Whether young Cahows always return to their exact point of departure, or whether they can return also to other islands/nesting colonies.

(9): Update on Oceanic Range Study of Cahow Using Archival Geolocational Data Loggers:

The 2010-2011 nesting season marked the second year that archival geolocational data loggers were deployed on the legs of selected adult Cahows. During the first season that these data loggers were deployed, a total of 10 of the 12 tags deployed were recovered successfully and the data they contained downloaded. Nine of the ten loggers recovered were found to contain useful data spanning periods ranging from 1 week to over 12 months. This data could then be converted into maps on which the daily positions of individual Cahows could be illustrated. After downloading data, these loggers were reset and re-deployed on the same or different birds to gain further locational data.

The data from these data loggers has completely revolutionized the understanding of the oceanic range of the Cahow, which is much more wide-ranging than originally thought. By combining all the useable data points from all nine of the recovered loggers, a new estimate for the overall range can now be shown (Fig. 9), covering an area stretching from the eastern coast of North America to Western European waters. In addition, 4 of the 6 loggers recovered with up to 12 months of data showed a distinct difference between their breeding and non-breeding ranges.




Fig. 9: Oceanic range of Cahow, combining all data points from 9 recovered loggers




Fig. 10: 12-month travels of one adult breeding Cahow as recorded by data logger

(10): Planned Future Management Actions and Research:

A number of the projects and proposals recommended in past yearly Nesting Season Reports have either been successfully completed or are well underway; following are the most important recommendations for continuing projects which are underway, or that are proposed for future nesting seasons:

2011 - 2012 Breeding Season:

- Start establishment efforts for second nesting colony on Nonsuch Island with planned translocation of 20 - 25 Cahow chicks from original nesting islets to “B” colony site on the South Hill of that island;
- Retrieve and download data from remaining Lotek LAT 2500 archival geolocational data loggers fitted to adult Cahows;
- Investigate DNA studies from blood sampling of Cahow to determine relationship with other North Atlantic Gadfly petrels, confirm that the Cahow is a distinct species, check for genetic diversity and variability within the population, check for possible inbreeding/lack of genetic variability and to confirm the sex of individual birds;
- Continue ongoing Cahow banding program;

- Continue use of sound attraction system at “A” translocation site on Nonsuch to continue attracting returning young Cahows to prospect for nests at that site;
- Continue monitoring nesting islands for presence of rats and other predators, and set out rodenticides when necessary.
- Continue installation of additional artificial nest burrows at nesting colonies

2012 – 2013 Breeding Season:

- Continue translocations of near-fledged Cahow chicks from nesting islands to “B” colony site on Nonsuch Island, moving 20 – 25 chicks annually until a target figure of 75 to 90 chicks have been moved and have fledged from Nonsuch;
- Install 2nd sound attraction system at new “B” colony site on Nonsuch Island and play disc of recorded Cahow courtship calls and cries during breeding season; also last year for using sound attraction system at original, “A” translocation site.
- Continue blood sampling of Cahows for DNA studies;
- Continue Cahow Banding program;
- Continue monitoring of nesting islands for the presence of rats; set out rodenticides when necessary;
- Continue installation of additional artificial nest burrows at nesting colonies.

(11). Acknowledgements:

A major species recovery project can not be successfully carried out without the presence of a large support group to provide manpower, backup and logistical support. The Terrestrial Conservation Officer would like to acknowledge with thanks the following Departmental staff, volunteers, organizations, schools and members of the public for their assistance in the Cahow Recovery Project during the 2010 – 2011 breeding season:

Mr. Drew Pettit, Director, Dept. of Conservation Services, Mr. Peter Drew, assistant Conservation Officer (Dept. of Conservation Services); Ms. Mandy Shailer (GPS co-ordinator, Dept. of Conservation Services), who for a second year has compiled locational data from the geolocational data loggers and used this to make the tracking and oceanic range maps, Nicholas Carlile and Dr. David Priddell, (New South Wales Department of the Environment and Climate Change, Australia), who have provided guidance on various aspects of the recovery program and are collaborating in the writing of two scientific papers on the breeding biology of the Cahow and the Cahow translocation project; Chris Flook, collections officer, Bermuda Aquarium, Museum and Zoo, Lynn Thorne, tours director, Bermuda Zoological Society (Nonsuch tours and “Cahow encounters” for school and BZS tour groups); Ben Watson and JP Rouja of LookTV (cameraman and producer for the “Higher Ground” documentary on the Cahow Translocation Project).

As usual, I am deeply indebted to the Terrestrial Conservation Crew for their hard work in building artificial concrete nest burrows for the Cahows on both Nonsuch Island and the original nesting islets. This work has consisted of making and transporting literally thousands of pounds of concrete out to the isolated nesting islands, landing it in buckets in generally rough conditions, and carrying it to the top of the islands. They have also carried out topping and removal of Casuarina trees at the translocation sites on Nonsuch Island to improve the habitat for Cahows to recolonize these areas. The Terrestrial Conservation Crew consists of Barry Smith (Foreman), Hillgrove Iris, Kiwon Furbert and Marvin Jones.

A good deal of the research work carried out by the Cahow Recovery Program has been possible only because of donations by schools, businesses, organizations and members of the public. For example, the geolocational loggers which have revolutionized the understanding of the oceanic range of the Cahow were purchased entirely through public donations, notably by Saltus Cavendish School and Paget Primary School. Special thanks also go out to the Atlantic Conservation Partnership (A.C.P.) and the Ascendant Group of Companies, who have supported and coordinated donations for the Nonsuch Island Living Museum and the Cahow Recovery Project. We are deeply grateful for the interest and generosity shown by all of these and other donors.

In conclusion, I would especially like to acknowledge my wife, Leila Madeiros, and children Seth and Elizabeth, without whose support and patience I would not have been able to carry out this demanding and time – consuming project over the last 10 years.

Jeremy Madeiros,
Terrestrial Conservation Officer



Fig. 11: Adult Cahow in IPO D1 nest burrow with large egg (January 2011)